

aastex emulateappj5,natbib

aa

$h \text{ min}^m \text{ sec}^s \text{ deg}^\circ$

N5 λ 1240 Si4 λ 1394 and λ 1403 Si4 λ 1394 Si4 λ 1403 C4 λ 1549 He2 λ 1640 C3] λ 1909 [Ne4] λ 2424 Mg2 λ 2800 [N2] λ 6548 [N2] λ 6583 [S2] λ 6716 [S2] λ 6731 [O3] λ 4959 [O3] λ 5007

Ly α N5 Si4 C4 He2 C3] [Ne4] Mg2 [N2] H α [S2] H β [O3]

$W_{\lambda,rest}$ H α (n) H α (b) $- .2ex \sim - .2ex \sim \sigma_{sky}$

Accepted for publication in the March 2003 Astronomical Journal.

Dawson et al. A High-Redshift, Hard X-ray Emitting Spiral

document

Optical and Near-Infrared Spectroscopy of a High-Redshift, Hard X-ray Emitting Spiral Galaxy1

Steve Dawson2, Nate McCrady2, Daniel Stern3, Megan E. Eckart4, Hyron Spinrad2, Michael C. Liu5, and James R. Graham2

1 Based on observations made at the W.M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation.

2 Department of Astronomy, University of California at Berkeley, Mail Code 3411, Berkeley, CA 94720 USA; sdawson@astro.berkeley.edu, nate@astro.berkeley.edu, spinrad@astro.berkeley.edu, jrg@astro.berkeley.edu. ■

3 Jet Propulsion Laboratory, California Institute of Technology, Mail Stop 169-327, Pasadena, CA 91109 USA; stern@zwolfkinder.jpl.nasa.gov.

4 Division of Physics, Mathematics, and Astronomy, California Institute of Technology, Mail Stop 220-47, Pasadena, CA 91125 USA; eckart@srl.caltech.edu.

5 Currently Beatrice Watson Parrent Fellow, the Institute for Astronomy, University of Hawai'i, 2680 Woodlawn Drive, Honolulu, HI 96822 USA; mliu@ifA.Hawaii.Edu.

abstract We present optical and near-infrared Keck spectroscopy of CXOHDFN J123635.6+621424 (hereafter HDFX28), a hard X-ray source at a redshift of $z = 2.011$ in the flanking fields of the Hubble Deep Field-North (HDF-N). HDFX28 is a red source ($\mathcal{R} - K_s = 4.74$) with extended steep-spectrum ($\alpha_{1.4GHz}^{8.4GHz} > 0.87$) microjansky radio emission and significant emission (441 μ Jy) at 15 μ m. Accordingly, initial investigations prompted the interpretation that HDFX28 is powered by star formation. Subsequent *Chandra* imaging, however, revealed hard ($\Gamma = 0.30$) X-ray emission indicative of absorbed AGN activity, implying that HDFX28 is an obscured, Type II AGN. The optical and near-infrared spectra presented herein corroborate this result; the near-infrared emission lines cannot be powered by star formation alone, and the optical emission lines indicate a buried AGN. HDFX28 is identified with a face-on, moderately late-type spiral galaxy. Multi-wavelength morphological studies of the HDF-N have heretofore revealed no galaxies with any kind of recognizable spiral structure beyond $z > 2$. We present a quantitative analysis of the morphology of HDFX28, and we find the measures of central concentration and asymmetry to be indeed consistent with those expected for a rare high-redshift spiral galaxy.